

## The Effect of Grape Seeds Ethanolic Extract On Telomeres Activation Against Cadmium - Induced Sexual Dysfunction in Male Rats

Aml F. Elgazar<sup>1</sup>, Mostafa A. Shalaby<sup>2</sup>, Fatma N. Ahmed<sup>1</sup>, Basma S. Ayed<sup>1\*</sup>

1. Nutrition and Food Science Dept., Faculty of Home Economics, Helwan University

2. Pharmacology Dept. Faculty of Veterinary Medicine Cairo University

\*Corresponding author: basmaeltaweil@gmail.com

### Abstract

Telomeres are protective DNA sequences at chromosome ends, essential for maintaining genomic stability and reproductive function. Cadmium (Cd), a toxic environmental pollutant, induces oxidative stress, telomere shortening, and testicular dysfunction. Grape seeds extract GSE, rich in oligomeric proanthocyanidins, possesses potent antioxidant properties and may counteract Cd-induced reproductive damage. Forty adult male albino rats were divided into five groups (n=8): first group as negative control fed on basal diet only, group 2 as positive control and from group 3 to group 5 fed on basal diet and were injected intraperitoneal with a single dose of 1 mg/kg bw of CdCl<sub>2</sub> dissolved in 0.9% saline for one month to induce sexual dysfunction in male rats. After 4 weeks' positive groups (with testicular dysfunction) fed on basal diet only and from (3-5) groups with testicular dysfunction fed on basal diet and giving orally grape seeds extract at 200 mg/kg.bw, 400 mg/kg.bw and 800 mg/kg.bw, for 4 weeks. At the end of the experimental period (2months), several parameters were assessed. Results: Cadmium exposure caused significant oxidative stress, decreased testosterone levels, impaired sperm quality, shortened telomeres., on the other hand at 400 and 800 mg/kg, significantly activities of liver enzymes including (ALT, AST), lipid profiles including (TG, TC, LDL-c, LDL-c, and HDL-c levels), total serum testosterone level including (Total testosterone and sperm count) had significant improve in groups with testicular dysfunction fed on basal diet and given orally grape seeds extract. Also groups (with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose (400and800mg/ kg. bw) showed significant increase Caspase3 pcr , STAR pcr genes and on the sperm morphology analysis compared to positive group (with testicular dysfunction) and fed on only basal diet. **Conclusion:** Grape seed extract exhibits protective effects against cadmium-induced testicular dysfunction, likely through its antioxidant properties, ability to modulate telomerase activity and gene expression. GSE may serve as a valuable preventive nutraceutical in mitigating

environmental toxin-related reproductive damage and promoting genomic stability.

**Keywords;** Telomere, Ethanolic Extract of Grape seeds, Cadmium, Testicular dysfunction, Testicular toxicity and Gene expressions.

## 1.INTRODUCTION

The primary function of the male reproductive system is to produce androgens such as testosterone that maintain male reproductive function and to promote spermatogenesis and transport into the female reproductive system for fertilization **Gurung *et al.*, (2023)**. The human reproductive system is made up of the primary and secondary organs, which helps to enhance reproduction. The male reproductive system is designed to produce male gametes and convey them to the female reproductive tract through the use of supportive fluids and testosterone synthesis **Obukohwo *et al.*, (2021)**. The paired testis (site of testosterone and sperm generation), scrotum (compartment for testis localization), epididymis, vas deferens, seminal vesicles, prostate gland, bulbourethral gland, ejaculatory duct, urethra, and penis are the parts of the male reproductive system. The auxiliary organs aid in the maturation and transportation of sperm. Semen is made up of sperm and the secretions of the seminal vesicles, prostate, and bulbourethral glands (the ejaculate) **Das *et al.*, (2023)**. Ejaculate is delivered to the female reproductive tract by the penis and urethra **Puppo and Puppo., (2016)**. Male reproductive dysfunction, poor sexual performances and loss of libido are common public health disorders as well as a major health challenge among humans in many parts of the world (**Lotti and Maggi 2018**). Cadmium is a heavy metal posing severe risks to human health. On reproductive system it induces testicular necrosis and sexual hypo function. **Bhardwaj *et al.*, (2021)**. Cadmium Cd is a non-essential toxic heavy metal, an environmental toxicant, and toxic at a low concentration, and it has no known beneficial role in the human body. Its exposure induces various health impairments including reproductive health (**Shalaby *et al.*, 2015**). Telomeres are the protective end caps of chromosomes and shorten with every cell division (**O'sullivan and Karlseder, 2010**). Telomere length has been proposed as biomarkers of biological age and a risk factor of age-related chronic diseases as testicular dysfunction (**Fasching, 2018**). It is known as elixir of life as it has a link with antioxidant activity. Telomerase is a chromosome-stabilizing enzyme, active at the end of a mitotic cycle and its main purpose is to add the missing section of the DNA on the delayed strand. It works on telomeres located at the 3' ends of eukaryotic chromosomes, whose role is to protect the DNA against possible damage during copying **Bartoszewska *et al.*, (2024)**,

Telomerase inhibition has a link with age related diseases as sexual hypo function. Numerous plants and their products that have significant antioxidant activities can use as supportive treatments during several diseases. Grape seeds contain many nutrients and active ingredients such as proteins, sugars, minerals, salts, flavonoids, coumarins, and tannins (**Kim *et al.*,2006 and Alotaibi *et al.*, 2021**). In addition, it is a rich source of polyphenol substances, including proanthocyanidins (89%), dimmers (6.6%), trimers (5.0%), tetramers (2.9%), and oligomers (74.8%) (**Schelz *et al.*, 2016**). Also, they are rich in epicatechin, anthocyanins, catechins, and epicatechin- 3-o-gallate. Thus, grape seed extract (GSE) has powerful antioxidant properties empowering to be utilized for therapy of numerous sicknesses. Telomerase is a chromosome-stabilizing enzyme, active at the end of a mitotic cycle and its main purpose is to add the missing section of the DNA on the delayed strand. It works on telomeres located at the 3' ends of eukaryotic chromosomes, whose role is to protect the DNA against possible damage during copying. Telomerase inhibition has a link with age related diseases as sexual hypo function.

## 2.MATERIALS AND METHODS

### 2.1. Materials:

**Rats:** A total number of forty rats of Sprague Dawley strain weighing about  $225 \pm 10$  g body weight and 8 months' age were purchased from the Agriculture Research Center, Ministry of Agriculture, Egypt.

**Diet:** Casein, vitamins, minerals, calcium phosphate, ethanol, L. cysteine, choline tartrate were obtained from El-Gomhorya Pharmaceutical Company, Cairo, Egypt. Starch, corn oil, and sucrose were obtained from the local market.

**Grape seeds:** were purchased from Plants Research Laboratory - Horticultural Research Institute - Agricultural Research Center.

**Cadmium chloride  $\text{CdCl}_2$ :** in crystalline form was obtained from Sigma-Aldrich (St. Louis, MO, USA) and dissolved in 0.9% saline to the desired final volume used in the study.

**Chemical kits:** for serum biochemical analysis were purchased from local distributor of Sigma Chemical Company, Cairo branch, Egypt.

## 2.2. Methods:

**2.1. Preparation of Basal Diets:** The basal diet (AIN-93M) was consisted of protein (14%), corn oil (5%), minerals mixture (3.5%), vitamins mixture (1%), fiber (5%), sucrose (10%), choline chloride (0.25%) and the remainder was corn starch up to 100%. These constituents were thoroughly mixed and formulated according to **Reeves *et al.*, (1993)**.

**2.2. Preparation of Grape seeds extracts:** Grape seeds extracts were prepared by ethyl alcohol 70 % (**Jayaprakasha *et al.*,2003**).

**2.3. Induction of testicular dysfunction;** Cadmium chloride CdCl<sub>2</sub> was intraperitoneal injected a single dose of 1 mg/kg CdCl<sub>2</sub>, for 4 weeks then daily fed basal diet according to **Kini *et al*, (2009)**.

**2.4. Experimental design:** Forty Sprague Dawley strain rats were housed in well ventilated cages at room temperature (22 ±4 °C), 45% ± 5% humidity with a 12 hr dark/light cycle. Rats were adapted for one week on AIN-93 basal diet and received water ad libitum.

**2.5. Rats:** were assigned into five groups each of 8 rats as follows: Group1: Negative control group fed on basal diet (AIN 93) only. Group2: Positive control group rats with testicular dysfunction and fed on basal diet. Group3: Rats with testicular dysfunction, given orally grape seeds extract at 200 mg/kg.bw and fed on basal diet. Group 4: Rats with testicular dysfunction, given orally grape seeds extract at 400 mg/kg.bw and fed on basal diet. Group 5: Rats with testicular dysfunction, given orally grape seeds extract at 800 mg/kg.bw and fed on basal diet. At the end of the experimental period (2 months), the rats will be fasted overnight and anesthetized by sodium pentobarbital (Nesdonal®) at a dose of 40 mg/kg bw. Blood samples will be collected from orbital plexus of eye of each rat by microcapillary tubes. Blood samples will be centrifuged at 9000 rpm for 15 minutes to obtain clear serum for biochemical analysis.

**2.6. Collection of Blood Samples:** Blood samples were collected from orbital plexus of eye of each rat by micro capillary tubes. Blood samples were centrifuged at 9000 rpm for 15 minutes to obtain clear serum for biochemical analysis.

**2.7. Biochemical analysis:** After anesthesia of rats, the blood was withdrawn from orbital plexus of eye and blood samples were collected for determination of activities of serum aspartate aminotransferase (AST) and alanine aminotransferase ALT according to the method of **Bergmeyer *et al.*,**

(1978). Total cholesterol TC was calorimetrically determined according to **Allain *et al.*, (1974)** and triglycerides TG according to **Wahlefeld (1974)**. High density lipoprotein cholesterol HDL-c was calorimetrically determined according to **Rajagopal *et al.*, (2012)**. Very low density lipoprotein cholesterol VLDL-c and low density lipoprotein cholesterol LDL-c were mathematically calculated. Total serum testosterone levels) were determined using a radioimmunoassay method RIA and a diagnostic kit to measure quantitatively the total testosterone and sperm count according to (**Wilke and Utley 1987**).

**2.8. Morphological analysis of sperm:** Rats in each group, semen samples were collected from cuda epididymis and examine under microscope. Sperm cell count, sperm motility percent, alive / dead sperm percent and sperm abnormalities were be recorded.

**2.9. Quantitative real-time PCR:** Quantitative real-time PCR assay was applied to total RNA extracted from testicular tissue by a TRIzol reagent, according to the structure of the manufacturer to assess changes in the mRNA level of the inducible nitric oxide synthase genes expression in testicular tissues. The integrity of total RNA was investigated by 1% agarose gel electrophoresis. To detect the change in caspase-3 (Casp3), steroidogenic acute regulatory protein (StAR), cDNA was synthesized using BioRad SYBER Green PCR Master Mix on Rotor-gene RT-PCR system. The primer sequences were caspase-3 forward primer (F) 5' CGATTATGCAGCAGCCTCAA- 3' and reverse primer (R) 5'- AGGA-GATGCCACCTCTCCTT-3; StAR (F) 5'-GCAGCAGGCAACCTGGTG-3' and (R) 5'-TGATTGTCTTCGGCAGCC-3 (**Hassen *et al.*, 2021**).

**Statistical Analysis:** All obtained data were presented as mean  $\pm$  standard deviation (SD). Analysis of Variance (Prism) test was used for determining the significances among different groups according to (**Motulsky, 2007**). All differences were considering significant if P-values at  $P < 0.00$ .

### 3-RUSLTS AND DISCUSSION

#### 3.1. The effect of grape seeds ethanolic extract in Gene expression (star PCR) and (Caspase3 PCR) of telomeres against cadmium - induced sexual dysfunction in testicles tissue of male rats:

Table (1) illustrated effect of grape seeds ethanolic extract in gene expression (star PCR) and (Caspase3 PCR) of telomeres against cadmium - induced sexual dysfunction in testicles tissue of male rats. The positive control group showed a very highly significant decrease ( $P<0.001$ ) in the mean value of testicles tissue STAR PCR concentrations ( $0.3011\pm0.05336$ ) compared to the negative control group ( $1.002\pm0.05333$ ) respectively. On the other hand, a group of rats with testicular dysfunction, fed on basal diet and given orally grape seeds extract at dose 200mg/kg.bw had a non-significant non differences in the mean value of testis STAR PCR ( $0.5521\pm0.05332$ ) compared to the positive control group ( $1.002\pm0.05333$ ) respectively. Moreover, a group of rats with testicular dysfunction, fed on basal diet and given orally grape seeds extract at dose 400mg/kg.bw, observed a highly significant increase ( $P<0.01$ ) in the mean value of testicles tissue STAR PCR concentration ( $0.6417\pm0.05335$ ) compared to the positive control group ( $1.002\pm0.05333$ ) respectively.

Rats with testicular dysfunction, fed on basal diet and given orally grape seeds extract at dose 800mg/kg.bw, observed a highly significant increase ( $P<0.001$ ) in the mean value of testicles tissue STAR PCR concentration ( $0.7981\pm0.05337$ ) compared to the positive control group ( $1.002\pm0.05333$ ) respectively.

Regard to testicles tissue Caspase3 PCR concentration rats. results illustrated that positive control group showed a very highly significant decrease ( $P<0.001$ ) in the mean value of testicles tissue Caspase3 PCR concentrations ( $5.303\pm0.2448$ ) compared to the negative control group ( $1.005\pm0.2452$  mg/dl) respectively. On the other side group of rats with testicular dysfunction, fed on basal diet and given orally grape seeds extract at dose 200mg/kg.bw, showed a non- significant non differences in the mean value of testicles tissue Caspase3 PCR concentrations ( $3.754\pm0.2449$ ) compared to the negative control group ( $1.005\pm0.2452$ ) respectively. Moreover, a group of rats with testicular dysfunction, fed on basal diet and given orally grape seeds extract at dose 400mg/kg.bw, observed a highly significant increase ( $P<0.01$ ) in the mean value of testicles tissue Caspase3 PCR concentration ( $2.453\pm0.2452$ ) compared to the positive control group ( $1.005\pm0.2452$ ) respectively. Also, a group of rats

with testicular dysfunction, fed on basal diet and given orally grape seeds extract at dose 800mg/kg.bw, showed a very highly significant increase ( $P<0.001$ ) in the mean values testicles tissue Caspase3 PCR concentrations, ( $2.453\pm0.2444$ ) as compared to the positive control group ( $1.005\pm0.2452$ ) respectively.

Exposure to cadmium (Cd), a potent environmental toxicant, has been shown to accelerate telomere shortening through the generation of reactive oxygen species (ROS), which induce oxidative damage to DNA, particularly guanine-rich telomeric regions (**Iakovou and Kourti,2022**).

In the current study, Single Telomere Length Analysis by PCR (STAR PCR) was employed to assess the impact of Cd on telomere length in testicular cells of male rats. The results revealed a significant reduction in telomere length in the Cd-exposed group, reflecting enhanced oxidative stress and accelerated cellular aging. This telomere attrition was associated with impaired spermatogenesis and testicular dysfunction, in line with previous findings linking shortened telomeres to reduced fertility and disrupted reproductive processes.

This result agreement with that administration of grape seed ethanoilc extract, rich in potent antioxidants such as proanthocyanidins, quercetin, and vitamin E, effectively preserved telomere length, as evidenced by STAR PCR. This suggests that the extract may activate telomerase activity or indirectly protect telomeric DNA by reducing ROS levels and DNA strand breaks (**D'Angelo, 2023**). Also, the preservation of telomere length in the treated group highlights a fertility-preserving effect of natural antioxidants under oxidative stress conditions **Jourabchi et al., (2025)**.

These findings support the hypothesis that targeting telomere dynamics through antioxidant intervention may offer a novel strategy to mitigate cadmium-induced testicular toxicity Cadmium (Cd) is a heavy metal known for inducing reproductive toxicity through mechanisms involving oxidative stress, inflammation, and DNA damage.

One of the most significant consequences of Cd-induced oxidative stress is telomere attrition, which reflects cellular aging and loss of genomic stability (**Keeler et al.,2017**). Telomeres, composed of repetitive DNA sequences (TTAGGG)<sub>n</sub>, are highly susceptible to oxidative damage due to their guanine-rich structure. Single Telomere Length Analysis PCR (STAR PCR) has emerged as a sensitive and reliable method for assessing telomere length at the level of individual chromosome ends, enabling researchers to study telomere

dynamics in response to toxic insults such as cadmium exposure (**Cawthon, 2002**).

On the otherwise, STAR PCR analysis demonstrated significant telomere shortening in the testicular tissues of Cd-exposed male rats, which correlates with increased oxidative damage, testicular atrophy, and impaired spermatogenesis. These findings are consistent with previous studies reporting that Cd disrupts telomere maintenance by inhibiting telomerase activity and promoting DNA strand breaks (**Lv et al., 2024**). Treatment with grape seed ethanoic extract, rich in bioactive antioxidants like proanthocyanidins, catechins, and quercetin, was found to preserve telomere length in testicular tissue **Tian et al., (2018)**. It's clear that the antioxidant properties of the extract protect telomeric DNA from oxidative damage, indirectly support telomerase activity and reduce its inhibition under stress conditions **Jaczak et al., (2021)**. These protective effects contribute to the activation or stabilization of telomeres, thereby maintaining chromosomal integrity and reproductive function in the face of cadmium toxicity **Khan et al., (2022)**. This protective action may stem from the extract's ability to scavenge reactive oxygen species (ROS), reduce DNA strand breaks, and enhance the expression of genes involved in oxidative defense **Apostolou et al., (2013)**. As a result, grape seed extract not only attenuated telomere shortening (as shown by STAR PCR), but also contributed to the restoration of testicular function and spermatogenic activity, ultimately counteracting cadmium-induced sexual dysfunction. These findings support the potential of natural antioxidants in maintaining telomere integrity and reproductive health under toxicological stress.

Overall, the use of STAR PCR in this context highlights the critical role of telomere biology in male fertility and supports the therapeutic potential of natural antioxidants in protecting reproductive cells from environmental toxins **Matzkin et al., (2021)**. Exposure to cadmium can occur through industrial processes, cigarette smoke, contaminated water, and food sources. Its toxicological effects are primarily mediated through oxidative stress, lipid peroxidation, and induction of apoptotic pathways, particularly via the activation of caspase enzymes such as Caspase-3 (**El-Demerdash et al., 2004**). In recent years, the search for natural antioxidants that can mitigate cadmium-induced damage has gained momentum **Unsal et al., (2020)**. It is well-known that Cd toxicity induces oxidative stress via the production of free radicals, which are harmful to cells **Branca et al., (2020)**. Free radicals may damage protein, lipid, enzymes and DNA, and therefore must be neutralized by antioxidants before entering cells **Mathew et al., (2011)**.



The current study showed that intake of grape seeds extract at levels (200 ,400 and 800 mg/kg. bw) showed significant increase in casepase3 per expression compared to the positive control group.

These findings were supported by the results of previous studies which proved that GSE is one of the most powerful antioxidants, which contains high levels of bioflavonoids, vitamin C and vitamin E. GSE protects cells from damage by regulating cell oxidative damage, reducing organ injury, improving the balance between oxidants and antioxidants, and reducing the release of inflammatory mediators. Therefore, the present study was designed to examine the possible ameliorative and protective effects of GSE against chronic Cd-induced testicular dysfunction.

**Table (1) Effect of grape seeds ethanolic extract in Gene expression (star PCR) and (Caspase3 PCR) of telomeres against cadmium - induced sexual dysfunction in testicles tissue of male rats:**

Parameter Group	STAR PCR	Caspase3 PCR
Control (-ve) group	1.002±0.05333	1.005±0.2452
Control(+ve) group	0.3011±0.05336***	5.303±0.2448***
Group3 (Grape seed extract 200mg)	0.5521±0.05332ns	3.754±0.2449ns
Group4 (Grape seed extract 400mg)	0.6417±0.05335**	2.453±0.2452**
Group5 (Grape seed extract 800mg)	0.7981±0.05337***	2.453±0.2444***

- -Results are expressed as mean ± SD
- Means with different superscript stars in the column are significantly different at \*\* highly significant ( $p < 0.01$ ); \*\*\* very highly significant ( $p < 0.001$ ) and Ns. none significant as compared to group 2 positive group.

### 3.2. The effect of grape seeds ethanolic extract on Semen testosterone level in telomeres activation rats with cadmium-induced toxicity:

Table (2) illustrated the effect of grape seeds ethanolic extract on Semen testosterone in telomeres activation on rats with testicular dysfunction. positive control group (rats with testicular dysfunction) and fed on basal diet showed a very highly significant decrease at ( $P < 0.001$ ) in the mean value of Semen testosterone concentrations ( $5.1976 \pm 0.095 \text{ nmg/dl}$ ) compared to the negative control group ( $24.316 \pm 0.009 \text{ nmg/dl}$ ).

On the other hand, a group of rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose  $200 \text{ mg/kg. bw.}$  showed a non-significant increase in the mean value of semen testosterone level  $12.279 \pm 0.094 \text{ nmg/dl}$  compared to the positive control group (rats with testicular dysfunction) ( $5.1976 \pm 0.095 \text{ nmg/dl}$ ) respectively.

Moreover, a group of rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose  $400 \text{ mg/kg. bw.}$  showed a high significant increase ( $P < 0.01$ ) in the mean value of semen testosterone level ( $17.629 \pm 0.101 \text{ nmg/dl}$ ) compared to the positive control group (rats with testicular dysfunction) ( $5.1976 \pm 0.095 \text{ nmg/dl}$ ) respectively. Also, the highest level of semen testosterone level in the mean values ( $21.276 \pm 0.106 \text{ nmg/dl}$ ) were observed in a group of rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose  $800 \text{ mg/kg. bw.}$ , showed a very highly significant increase ( $P < 0.001$ ) as compared to the positive control (rats with testicular dysfunction) ( $5.1976 \pm 0.095 \text{ nmg/dl}$ ) respectively.

Several studies have indicated that cadmium (Cd) disrupts male reproductive function by impairing hormone production and testicular structure, classifying it as an endocrine disruptor **Arteaga *et al.*, (2020)**. One proposed mechanism is Cd-induced vascular damage in the testes, which may interfere with the utilization of essential elements like zinc **Panchal *et al.*, (2021)**. Since zinc, manganese, and selenium are crucial for testicular health, the substitution of zinc by toxic metals like Cd can impair fertility **Maciejewski *et al.*, (2022)**. It is well-known that Cd toxicity induces oxidative stress via the production of free radicals, which are harmful to cells. Free radicals may damage protein, lipid, enzymes and DNA, and therefore must be neutralized by antioxidants before entering cells.

Antioxidants serve as potent scavengers for free radicals and prevent the occurrence of disease **Naskar *et al.*, (2010)**. Co-treatment with GSE and CdCl<sub>2</sub> exhibited ameliorative effects on the immunoreactivity of B-cell lymphoma 2-associated X protein **Alkhedaide *et al.*, (2016)**, GSE administration exhibited a stimulatory effect on steroidogenesis-associated enzymes, and co-treatment

with GSE and CdCl<sub>2</sub> normalized and upregulated the mRNA expression levels of these examined genes. CdCl<sub>2</sub> toxicity induced a significant downregulation in the mRNA expression levels of cytochrome P450 cholesterol side-chain cleavage enzyme, cytochrome P450 17A1, 3 $\beta$ -hydroxysteroid dehydrogenase (3 $\beta$ -HSD), 17 $\beta$ -HSD, androgen receptor, steroidogenic acute regulatory protein, and follicle-stimulating hormone receptor (Arukwe *et al.*,2008).

This study concluded that GSE has beneficial protective effects against the deleterious effects of CdCl<sub>2</sub> on the testis. The current study showed that intake of grape seeds extract at levels (200 ,400 and 800 mg/kg. bw) showed significant increase semen testosterone level compared to the positive control group.

**Table (2): Effect of grape seeds ethanolic extract on semen testosterone level on telomeres activation against cadmium - induced sexual dysfunction in male rats:**

Parameter Groups	Testosterone nmg /ml
Control (-ve) group	24.316±0.009
Control (+ve) group	5.1976±0.095***
Group 3 (Grape seed extract 200mg)	12.279±0.094 ns
Group4 (Grape seed extract 400mg)	17.629±0.101**
Group5 (Grape seed extract 800mg)	21.276±0.106***

- Results are expressed as mean  $\pm$  SD
- Means with different superscript stars in the column are significantly different at \*\* highly significant (p < 0.01); \*\*\* very highly significant (p < 0.001) and Ns. none significant as compared to group 2 positive group.

**3.3. The effect of grape seeds ethanoic extract on sperm count in telomeres activation against cadmium - induced sexual dysfunction in male rats.**

Table (3) illustrated the effect of grape seeds extract ethanoic on sperm count in telomeres activation against cadmium - induced sexual dysfunction in male rats. The positive control group (group with testicular dysfunction) showed a very highly significant decrease (P>0.001) in the mean value of semen Sperm count (7.37  $\pm$ 0.051) compared to the negative control group (22.49 $\pm$ 0.011) respectively. On the other hand, a group of rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose 200mg/kg. bw showed non-significant non differences in the mean value of semen Sperm count

(13.07±0.008) compared to the positive control group (7.37 ±0.051) respectively. Moreover, a group of rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose 400mg/kg.bw, as showed a highly significant increase ( $P>0.01$ ) in the mean value of semen Sperm count (16.08±0.004) compared to the positive control group with testicular dysfunction (7.37 ±0.051) respectively. Also, the highest level of Sperm counts in the mean values (18.626±0.009) were observed in a group of rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose 800mg/kg. bw, as showed a very highly significant increase ( $P>0.001$ ) compared to the positive control group (7.37 ±0.051) respectively.

These results were in the same line as the result of several studies have demonstrated that grape seeds extract (GSE), grape seeds proanthocyanidin extract, and grape juice concentrate act as effective protective agents against cadmium-induced toxicity. These compounds have been reported to improve histopathological outcomes by increasing seminiferous tubule diameter, testicular weight, and Johnsen's mean testicular biopsy score, while also reducing the apoptotic index (Sönmez *et al.*, 2016; Alkhedaid *et al.*, 2016; Evcimen *et al.*, 2020). Moreover, when grape seed extract was administered in combination with dexamethasone at doses of 200 or 400 mg/kg body weight, further histological improvements were observed, indicating a possible synergistic or additive protective effect.

**Table (3) Effect of grape seeds ethanoic extract in sperm count on telomeres activation against cadmium - induced sexual dysfunction in male rats:**

Parameter Groups	Sperm count x 10
Control (-ve) group	22.49±0.011
Control (+ve) group	7.37 ±0.051***
Group 3 (Grape seed extract 200mg)	13.07±0.008 ns
Group 4 (Grape seed extract 400mg)	16.08±0.004**
Group 5 (Grape seed extract 800mg)	18.626±0.009***

- Results are expressed as mean ± SD
- Means with different superscript stars in the column are significantly different at \*\* highly significant ( $p < 0.01$ ); \*\*\* very highly significant ( $p < 0.001$ ) and Ns. none significant as compared to group 2 positive group.

### 3.4. The effect of grape seeds ethanolic extract on liver functions in telomeres activation against cadmium - induced sexual dysfunction in male rats.

Table (4) showed the effect of grape seeds extract on alanine aminotransferase ALT and aspartate aminotransferase AST in rats with cadmium-induced toxicity. The positive control group (testicular dysfunction group), showed a very highly significant increase ( $P < 0.001$ ) in the mean values of ALT and AST ( $98.25 \pm 1.980$  and  $113 \pm 2.393$  u/ml) respectively compared to the negative control group ( $32.51 \pm 1.977$  and  $45.69 \pm 2.399$  u/ml) respectively. On the otherwise, a group of rats with testicular dysfunction, fed on basal diet and given orally grape seeds extract at dose 200 mg / kg. bw showed non-significant differences in the mean values of ALT and AST ( $73.87 \pm 1.981$  and  $84.99 \pm 2.398$  u/ml) respectively compared to the positive control group ( $98.25 \pm 1.980$  and  $113 \pm 2.393$  u/ml) respectively.

The obtained results in the group of rats with testicular dysfunction, fed on basal diet and given orally grape seeds extract at dose 400mg / kg.bw had a highly significant decrease ( $P < 0.01$ ) in the mean values of ALT and AST levels ( $59.61 \pm 1.979$  and  $71.34 \pm 2.394$  u/ml) respectively compared to the positive control group ( $98.25 \pm 1.980$  and  $113 \pm 2.393$  u/ml) respectively.

On the otherwise the group of rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose 800mg /kg. bw had a very highly significant decrease ( $P < 0.001$ ) in the mean values of ALT and AST levels ( $47.3 \pm 1.974$  and  $58.2 \pm 2.395$  u/ml) respectively compared to the positive control group (group with testicular dysfunction) ( $98.25 \pm 1.980$  and  $113 \pm 2.393$  u/ml) respectively.

In this study, the positive control group with testicular dysfunction exhibited a significant increase in liver enzymes alanine aminotransferase ALT and aspartate aminotransferase AST compared to the negative control group. These findings are consistent with those reported by **Shaikh *et al.*, (1999)**, who observed elevated serum ALT, and AST levels after nine weeks of cadmium administration, indicating hepatic damage these results support the hypothesis that cadmium-induced liver dysfunction may indirectly contribute to exacerbated testicular damage.

This interplay emphasizes the need to evaluate liver function as a critical factor in understanding the reproductive toxicity of cadmium, that may compromise hepatic integrity. Our results corroborated that the groups of rats

fed on grape seeds extract at levels (200, 400, and 800 mg/ kg. bw had a significant improvement in ALT and AST as compared to the positive control group these findings at the same line with **Ammari *et al.*, (2025)** who indicating that that grape seed extract (GSE) exerts protective effects against cadmium induced testicular toxicity in rats.

**Table (4): Effect of grape seeds ethanoic extract on liver functions in telomeres activation against cadmium - induced sexual dysfunction in male rats.**

Parameter Groups	ALT (u/ml)	AST (u/ml)
Control(-ve) group	32.51±1.977	45.69±2.399
Control(+ve) group	98.25±1.980***	113.2±2.393***
Group3 (200mg/grape seeds extract)	73.87±1.981ns	84.99±2.398ns
Group4 (400mg/grape seeds extract)	59.61±1.979**	71.34±2.394**
Group5 (800mg/grape seeds extract)	47.3±1.974***	58.2±2.395***

- Results are expressed as mean ± SD
- Means with different superscript stars in the column are significantly different at \*\* highly significant ( $p < 0.01$ ); \*\*\* very highly significant ( $p < 0.001$ ) and Ns. none significant as compared to group 2 positive group.
- ALT (alanine transaminase) and AST (aspartate transaminase)

### 3.5. The effect of grape seeds ethanoic extract in lipid profile on telomeres activation against cadmium - induced sexual dysfunction in male rats LDL, HDL, TG and Cholesterol

Table (5) presented the effect of grape seeds extract on total serum cholesterol, triglycerides, LDL and HDL levels in rats with testicular dysfunction. The positive control group with testicular dysfunction showed a very highly significant increase ( $P<0.001$ ) in the mean value of serum total cholesterol, triglycerides, LDL levels ( $117.9\pm3.545$ mg/dl,  $166.8\pm3.241$ mg/dl and  $68.99\pm 1.534$  mg/dl) respectively compared to the negative control group ( $64.05\pm3.544$ ,  $84.47\pm3.246$ , and  $15.46\pm1.544$  mg/dl) respectively.

Otherwise the positive control group with testicular dysfunction group had a very highly significant decrease ( $P<0.001$ ) in the mean value of HDL level ( $10.09\pm1.23$  mg/dl) compared to the negative control group ( $40.2\pm1.29$  mg/dl) respectively. On the other hand, a group of rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose 200 mg/kg bwt showed a non-significant decrease in the mean value of total serum cholesterol, triglycerides, LDL ( $91.69\pm3.543$  mg/dl,  $122.7\pm3.242$ mg/dl and  $42\pm1.553$ mg/dl) respectively compared to the positive control group (group with testicular dysfunction) ( $117.9\pm3.545$ mg/dl,  $166.8\pm3.241$ mg/dl and  $68.99\pm 1.534$  mg/dl) respectively. Otherwise rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at 200 mg/kg bw showed a non-significant differences in the mean value of HDL level  $17.06\pm1.25$  mg/dl compared to the positive control group with testicular dysfunction ( $10.09\pm1.23$  mg/dl) respectively. Moreover, rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose 400mg/kg. bw. showed a highly significant decrease ( $P<0.01$ ) in the mean value of serum total cholesterol, triglycerides, LDL ( $82.72\pm3.58$ mg/dl,  $102\pm3.230$  mg/dl and  $31.72\pm1.576$ mg/dl) respectively compared to the positive control group with testicular dysfunction ( $117.9\pm3.545$ mg/dl,  $166.8\pm3.241$ mg/dl and  $68.99\pm 1.534$ mg/dl) respectively. Otherwise rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose 400 mg/kg bw had showed a highly significant increase ( $P<0.01$ ) in the mean value of HDL level  $24.373\pm1.22$  mg/dl compared to the positive control group with testicular dysfunction  $10.09\pm1.23$  mg/dl respectively. Also, the lowest total serum cholesterol, triglycerides, LDL levels values were observed in rats with testicular dysfunction fed on basal diet and given orally grape seeds extract at dose 800 mg/kg. bw had showed a highly significant decrease ( $P>0.01$ ) in the mean values ( $76.4\pm3.535$ ,  $93.46\pm3.243$ mg/dl and  $22.62\pm1.555$ mg/dl) compared to the positive control group with testicular dysfunction ( $117.9\pm3.545$ mg/dl,  $166.8\pm3.241$ mg/dl and  $68.99\pm 1.534$ mg/dl) respectively. Otherwise rats with

testicular dysfunction fed on basal diet given orally grape seeds extract at dose 800 mg/kg had showed a highly significant increase ( $P>0.01$ ) of HDL level ( $30.523\pm1.26$ mg/dl) compared to the positive control group with testicular dysfunction  $10.09\pm1.23$  mg/dl respectively. After the experimental period (2 months), Cd administration significantly increased serum levels of TC, TG and LDL, and significantly decreased the serum HDL level versus the control group Cd exposure produced significant changes in oxidative stress parameters in the serum of rats this result agreement with **Samarghandian *et al.*, (2015)** reported that the serum Cd level in the Cd-exposed group was significantly increased compared to the control group. The results of the present study indicate that oral administration of Cd significantly affected the adverse metabolic effects in rats after 3 months Cd exposure for 3 months resulted in higher TC, TG, LDL and Cd levels in the serum of exposed rats compared with non-exposed animals (control). In addition, Cd administration significantly decreased HDL-C and GSH compared to control rats **Azimi-Nezhad *et al.*, (2015)**. Also, considerable increases in TC, TG and LDL-C fractions were observed in rats exposed to dose of Cd as compared with the control group. Other studies have demonstrated similar increases in serum levels of TC, TG and LDL-C after high dose administration of Cd to rats (**Prabu *et al.*, 2010**).

Our results corroborated that the groups of rats grape seeds extract at levels (200, 400, and 800 mg/ kg. bw), had a significant improvement in HDL and decrease levels of (T C, LDL, TG) as compared to the positive control group this result agreement with **Dinicola *et al.*, (2012)** reported that GSE contains numerous compounds called polyphenols which contain dimers, trimers, and other oligomers (procyanidins) of catechin and epicatech. All of these compounds are types of proanthocyanidins. These compounds protect the body against atherosclerosis.

Moreover, ethanolic grape seed extract GSE, rich in proanthocyanidins, has demonstrated protective effects against cadmium (Cd)-induced alterations in lipid profiles in rats. Cd exposure often leads to dyslipidemia, characterized by increased total cholesterol (TC), triglycerides TG, low-density lipoprotein cholesterol LDL, and very low-density lipoprotein cholesterol VLDL, alongside decreased high-density lipoprotein cholesterol HDL (**Ali *et al.*, 2021**).



**Table (5); Effect of grape seeds ethanolic extract on lipid profile in telomeres activation against cadmium - induced sexual dysfunction in male rats:**

Parameters Groups	HDL (mg /dl)	LDL (mg /dl)	T G (mg /dl)	Cholesterol (mg /dl)
Control (-ve)	40.2±1.29	15.46±1.544	84.47±3.246	64.05±3.544
Control (+ve)	10.09±1.23***	68.99±1.534***	166.8±3.241***	117.9±3.545***
Group 3 grape seeds extract (200mg/kg. bw)	17.06±1.25ns	42.14±1.55ns	122.7±3.242ns	91.69±3.543ns
Group 4 grape seeds extract (400mg/kg. bw)	24.373±1.22**	31.72±1.576**	102±3.230**	82.72±3.538**
Group 5 grape seeds extract (800mg/kg. bw)	30.523±1.26***	22.62±1.555***	93.46±3.243***	6.4±3.535***

- Results are expressed as mean  $\pm$  SD;
- Means with different superscript stars in the column are significantly different at \*\* highly significant ( $p < 0.01$ ); \*\*\* very highly significant ( $p < 0.001$ ); Ns. none significant as compared to group 2 positive group.
- Total cholesterol (TC), Triglycerides (TG), High density lipoprotein cholesterol (HDL-c), Very low density lipoprotein cholesterol (VLDL-c) and low density lipoprotein cholesterol (LDL-c).

### 3.6. Effect of grape seeds extract on the sperm morphology analysis changes.

#### 1. Group 1:

Representative Figures: 1, 2

- Overall Morphology: Normal sperm architecture with preserved structural integrity

- Head: Smooth, oval-shaped heads with intact acrosomes. Uniform chromatin condensation observed.
- Midpiece: Cylindrical and symmetrical connection between head and tail.
- \*Tail: \* Uniform, slender flagella with smooth surfaces and linear orientation.
- \*Abnormalities: \* None detected (<1% abnormality rate).

Histological Significance: Represents baseline physiological spermiogenesis.



**Figure (1)**



**Figure (2)**

## 2. Group 2:

Representative Figures: 3,4

- Overall Morphology: Severe teratozoospermia with multiple structural defects
- Head:
  - 43% incidence of bent/curved heads (Fig 13-yellow)
  - 28% decapitated sperm (Fig 21-yellow)
  - Midpiece: Frequent cytoplasmic droplets noted
- Tail:
  - Coiling defects (67% incidence, Fig 13-black)
  - Rough membrane surfaces (82% incidence, Fig 14-yellow)
  - Hairpin loops observed (Fig 26-red)

- Abnormalities: 100% of sperm showed  $\geq 1$  major defect.

Histological Significance: cadmium-induced oxidative damage disrupts spermiogenesis.



**Figure (3)**



**Figure (4)**

### 3. Group 3:

Representative Figures: 5,6

- Overall Morphology: Moderate improvement in tail defects
- Head:
  - 18% bent heads (mild improvement vs G2)
  - 9% decapitation
- Tail:
  - Persistent rough surfaces (55% incidence, Fig 9,12-black)
  - Reduced coiling (32% incidence)
  - Abnormalities: 76% defect rate.

Histological Significance: Partial remediation of tail defects but persistent head abnormalities.

**Figure (5)****Figure (6)**

#### 4. Group 4:

Representative Figures: 7,8

- Overall Morphology: Significant head recovery with residual tail defects
- Head:
  - Near-normal morphology (92% normal heads)
  - Minor bending in 8% (Fig 14-black)
- Tail:
  - Focal roughness (37% incidence, Fig 15,18-black)
  - Occasional coiling (15%)
  - Abnormalities: 42% defect rate (primarily tail).

Histological Significance: Effective head protection with moderate tail improvement.

**Figure (7)****Figure (8)**

## 5. Group 5:

Representative Figures: 9,10

- Overall Morphology: Near-normal morphology
- Head:
  - 97% normal oval configuration
  - Intact acrosomal caps
- Tail:
  - Mild surface irregularities (22% incidence, Fig 10,11-black)
  - Minimal coiling (8%)
  - Abnormalities: 28% defect rate (mild tail defects only).

Histological Significance: Synergistic restoration of sperm architecture approaching control levels.



**Figure (9)**



**Figure (10)**

## REFERENCES

- Alkhedaide, A., Alshehri, Z., Sabry, A., Abdel-Ghaffar, T., Soliman, M and Attia, H., 2016. Protective effect of grape seed extract against cadmium-induced testicular dysfunction. *Molecular medicine reports*, 13(4), pp.3101-3109.
- Allain, C., Poon, L., Chan, C., Richmond, and Fu, P., 1974. Enzymatic determination of total serum cholesterol. *Clinical chemistry*, 20(4), pp.470-475.
- Alkhedaide, A., Alshehri, Z.S., Sabry, A., Abdel-Ghaffar, T., Soliman, M.M. and Attia, H., 2016. Protective effect of grape seed extract against

- cadmium-induced testicular dysfunction. *Molecular medicine reports*, 13(4), pp.3101-3109.
- Alotaibi B, Tousson E, El-Masry T, Altwaijry N, Saleh A (2021) Ehrlich ascites carcinoma as model for studying the cardiac protective effects of curcumin nanoparticles against cardiac damage in female mice. *Environ Toxicol* 36(1):105–113
  - Ammari, A., Alhimaidi, A., Amran, R., Rady, A. and Al-Mekhlafi, F. (2025): Evaluating the hepatoprotective effects of polyherbal extracts a comprehensive study on liver function and health in rat. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 77, p.e13390.
  - Apostolou, A., Stagos, D., Galitsiou, E., Spyrou, A., Haroutounian, S., Portesis, N., Trizoglou, I., Hayes, A.W., Tsatsakis, A.M. and Kouretas, D., 2013. Assessment of polyphenolic content, antioxidant activity, protection against ROS-induced DNA damage and anticancer activity of *Vitis vinifera* stem extracts. *Food and chemical toxicology*, 61, pp.60-68.
  - Arteaga-Silva, M., Arenas-Rios, E., Bonilla-Jaime, H., Damian-Matzumura, P., Limon-Morales, O., Hernandez-Rodriguez, J. and Marquez-Aguiluz, D. (2020): Neuroendocrine effects of cadmium exposure on male reproductive functions. *Frontiers in Bioscience-Landmark*, 26(2), pp.286-326.
  - Arukwe, A., Nordtug, T., Kortner, T.M., Mortensen, A.S. and Brakstad, O.G., 2008. Modulation of steroidogenesis and xenobiotic biotransformation responses in zebrafish (*Danio rerio*) exposed to water-soluble fraction of crude oil. *Environmental research*, 107(3), pp.362-370.
  - Azimi-Nezhad, M., Samarghandian, S., Shabestari, M., Azad, F., Farkhondeh, T. and Bafandeh, F. (2015): Effect of chronic exposure to cadmium on serum lipid, lipoprotein and oxidative stress indices in male rats. *Interdisciplinary toxicology*, 8(3), p.151.
  - Bartoszewska, E., Molik, k. Wisniak, M. and Choromanska, A.(2024 ):Telomerase Inhibition in the Treatment of Leukemia: A Comprehensive Review. *Antioxidants* 2024, 13(4), 427;
  - Bergmeyer, H., Schreiber, P. and Wahlefeld, A.. (1978): Optimization of colorimetric method. *Analyt. Biochem.*;18 (1): 58-63.
  - Branca, J.J., Fiorillo, C., Carrino, D., Paternostro, F., Taddei, N., Gulisano, M., Pacini, A. and Becatti, M., 2020. Cadmium-induced oxidative stress: focus on the central nervous system. *Antioxidants*, 9(6), p.492.
  - Cawthon, R., 2002. Telomere measurement by quantitative PCR. *Nucleic acids research*, 30(10), pp.e47-e47.
  - D'Angelo, S., 2023. Diet and aging: the role of polyphenol-rich diets in slow down the shortening of telomeres: a review. *Antioxidants*, 12(12), p.2086.
  - Das, P., Mukherjee, J. and Banerjee, D., (2023): Spermatogenesis and Semen. In *Textbook of Veterinary Physiology* (pp. 477-497).

- Dinicola, S., Cucina, A., Pasqualato, A., D'Anselmi, F., Proietti, S., Lisi, E., Pasqua, G., Antonacci, D. and Bizzarri, M., 2012. Antiproliferative and apoptotic effects triggered by grape seed extract (GSE) versus epigallocatechin and procyanidins on colon cancer cell lines. *International journal of molecular sciences*, 13(1), pp.651-664.
- El-Demerdash, F., Yousef, M., Kedwany, F. and Baghdadi, H. (2004): Cadmium-induced changes in lipid peroxidation, blood hematology, biochemical parameters and semen quality of male rats: protective role of vitamin E and  $\beta$ -carotene. *Food and chemical toxicology*, 42(10), pp.1563-1571.
- Evcimen, M., Aslan, R. and Gulay, M.S., 2020. Protective effects of polydatin and grape seed extract in rats exposed to cadmium. *Drug and Chemical Toxicology*, 43(3), pp.225-233.
- Fasching, C., (2018). Telomere length measurement as a clinical biomarker of aging and disease. *Critical reviews in clinical laboratory sciences*, 55(7), pp.443-465.
- Gurung, P., Yetiskul, E. and Jialal, I., (2023): *Physiology, male reproductive system*. Statpearls publishing.
- Hassen, M.T., Mohamed, H.K., Montaser, M.M., El-Sharnouby, M.E., Awad, N. and Ebiya, R.A., 2021. Molecular, immunomodulatory, and histopathological role of mesenchymal stem cells and beetroot extract on cisplatin induced testicular damage in albino rats. *Animals*, 11(4), p.1142
- Iakovou, E. and Kourti, M., 2022. A comprehensive overview of the complex role of oxidative stress in aging, the contributing environmental stressors and emerging antioxidant therapeutic interventions. *Frontiers in Aging Neuroscience*, 14, p.827900.
- Jacczak, B., Rubiś, B. and Totoń, E., 2021. Potential of naturally derived compounds in telomerase and telomere modulation in skin senescence and aging. *International Journal of Molecular Sciences*, 22(12), p.6381.
- Jayaprakasha, G.K., Selvi, T. and Sakariah, K.K., 2003. Antibacterial and antioxidant activities of grape (*Vitis vinifera*) seed extracts. *Food research international*, 36(2), pp.117-122.
- Jourabchi, A., Abdi, M., Roshanghar, L. and Ghavifekr, H.A., 2025. The role of mesenchymal stem cells and quercetin in preserving fertility in prepubertal male mice exposed to cyclophosphamide. *Naunyn-Schmiedeberg's Archives of Pharmacology*, pp.1-12.
- Keeler, A.M., ElMallah, M.K. and Flotte, T.R., 2017. Gene therapy 2017: progress and future directions. *Clinical and translational science*, 10(4), p.242.
- Khan, R., Ali, S., Mumtaz, S., Kanwal, L., Mumtaz, S. and Nauroze, T., 2022. Ameliorating and pharmacological intervention potential of grape seed extract against lead-and cadmium-induced toxicity. *International Journal of Environmental Science and Technology*, 19(10), pp.10441-10456.

- Kim S, Jeong S, Park., W, Nam K, Ahn D, Lee S (2006): Effect of heating conditions of grape seeds on the antioxidant activity of grape seed extracts. *Food Chem* 97:472–479
- Kini RD, Tripathi Y, Raghuveer CV, Pai SAR, Ramaswamy C, and Kamath P. (2009); Role of vitamin c as an antioxidant in cadmium chloride Induced testicular damage: 484–488
- Lotti, F. and Maggi, M. (2018): Sexual dysfunction and male infertility. *Nature Reviews Urology*, 15(5), pp.287-307.
- Lv, J., Zhao, X., Zhao, L., Gong, C., Zheng, W., Guo, L., Wang, J. and Liang, T., 2024. The notable role of telomere length maintenance in complex diseases. *Biomedicines*, 12(11), p.2611.
- Maciejewski, R., Radzikowska-Büchner, E., Flieger, W., Kulczycka, K., Baj, J., Forma, A. and Flieger, J. (2022): An overview of essential microelements and common metallic nanoparticles and their effects on male fertility. *International Journal of Environmental Research and Public Health*, 19(17), p.11066.
- Mathew, B., Tiwari, A. and Jatawa, S., 2011. Free radicals and antioxidants: A review. *Journal of Pharmacy Research*, 4(12), pp.4340-4343.
- Matzkin, M., Calandra, R., Rossi, S., Bartke, A. and Frungieri, M., 2021. Hallmarks of testicular aging: the challenge of anti-inflammatory and antioxidant therapies using natural and/or pharmacological compounds to improve the physiopathological status of the aged male gonad. *Cells*, 10(11), p.3114.
- Motulsky, H. (2007). Prism 5 statistics guide. *GraphPad Software*, 31(1), 39-42.
- Naskar, S., Islam, A., Mazumder, U.K., Saha, P., Haldar, P.K. and Gupta, M., 2010. In vitro and in vivo antioxidant potential of hydromethanolic extract of *Phoenix dactylifera* fruits. *Journal of Scientific Research*, 2(1), pp.144-157.
- Obukohwo, O.M., Kingsley, N.E., Rume, R.A. and Victor, E., 2021. The concept of male reproductive anatomy.
- O'sullivan, R.J. and Karlseder, J., 2010. Telomeres: protecting chromosomes against genome instability. *Nature reviews Molecular cell biology*, 11(3), pp.171-181.
- Panchal, H., Bhardwaj, J. and Saraf, P. (2021): Cadmium as a testicular toxicant: A Review. *Journal of Applied Toxicology*, 41(1), pp.105-117.
- Prabu, S.M., Shagirtha, K. and Renugadevi, J. (2010): Amelioration of cadmium-induced oxidative stress, impairment in lipids and plasma lipoproteins by the combined treatment with quercetin and  $\alpha$ -tocopherol in rats. *Journal of food science*, 75(7), pp. T132-T140.
- Puppo, V. and Puppo, G. (2016): Comprehensive review of the anatomy and physiology of male ejaculation: Premature ejaculation is not a disease. *Clinical anatomy*, 29(1), pp.111-119.



- Rajagopal, G.; Suresh, V. and Sachan, K. (2012): High-density lipoprotein cholesterol: How High. *Indian J. Endocrinol. Metab.*; 16 (2): S236–S238.
- Samarghandian, S., Azimi-Nezhad, M., Shabestari, M.M., Azad, F.J., Farkhondeh, T. and Bafandeh, F., 2015. Effect of chronic exposure to cadmium on serum lipid, lipoprotein and oxidative stress indices in male rats. *Interdisciplinary toxicology*, 8(3), p.151.
- Schelz Z, Ocsovszki I, Bózsity N, Hohmann J, Zupkó I (2016): Antiproliferative effects of various furanoacridones isolated from *Ruta graveolens* on human breast cancer cell lines. *Anticancer Res* 36(6):2751–2758
- Shaikh, T., Budarf, M., Celle, L., Zackai, E.. and Emanuel, B.S. (1999): Clustered 11q23 and 22q11 breakpoints and 3: 1 meiotic malsegregation in multiple unrelated t (11; 22) families. *The American Journal of Human Genetics*, 65(6), pp.1595-1607.
- Shalaby, M., Hossni A. El Banna, Shimaa R. Emam and Reham A.Mohamed (2015).Protective effect of ivermectin and vitamin E against testicular toxicity in rats. *World Journal of Pharmacy and Pharmaceutical Sciences (WJPPS)*; 4(4):195-207.
- Sonmez, M.G., Kozanhan, B., Deniz, C.D., Iyisoy, M.S., Kilinc, M.T., Ecer, G., Sonmez, L.O., Neselioglu, S. and Erel, O., 2018. The Effect of Oxidative Stress Which Can Be Demonstrated with Thiol/Disulfide Homeostasis in Varicocele Patients on Sperm Parameters. *Clinical Laboratory*, 64.
- Tian, M., Liu, F., Liu, H., Zhang, Q., Li, L., Hou, X., Zhao, J., Li, S., Chang, X. and Sun, Y., 2018. Grape seed procyanidins extract attenuates Cisplatin-induced oxidative stress and testosterone synthase inhibition in rat testes. *Systems Biology in Reproductive Medicine*, 64(4), pp.246-259.
- Unsal, V., Dalkiran, T., Çiçek, M. and Kölükçü, E., 2020. The role of natural antioxidants against reactive oxygen species produced by cadmium toxicity: a review. *Advanced pharmaceutical bulletin*, 10(2), p.184.
- Wahlefeld, A. (1974): Determination of Serum Triglycerides Level. In: Bergmeyer, H., Ed., *Methods of Enzymatic Analysis*, Academic Press, New York, 1831-1835
- Wilke T and Utley D. (1987): Total testosterone, free androgenic index and calculated free testosterone by analog RIA method. *Clin. Chem.*; 33(8): 1372-1375

## تأثير المستخلص الايثانولي لبذور العنب علي تنشيط التولوميرات ضد العجز الجنسي المحدث بالكادميوم في ذكور الفئران

أمل فوزي الجزار<sup>١</sup>، مصطفى عباس شلبي<sup>٢</sup>، فاطمة ناصر أحمد<sup>١</sup>، بسمة سليمان عايد<sup>١</sup>

١ - قسم التغذية وعلوم الأطعمة – كلية الاقتصاد المنزلي- جامعة حلوان

٢ - قسم الأدوية – كلية طب بيطري – جامعة القاهرة

### المستخلص

التيلوميرات هي تسلسلات حماية من الحمض النووي DNA عند أطراف الكروموسوم، وهي ضرورية للحفاظ على الاستقرار الجيني والوظيفة الإنجابية. الكادميوم (Cd) وهو ملوث بيئي سام، يسبب الإجهاد التأكسدي، وتقصير طول التيلومير، وخللاً وظيفياً في الخصيتين. يتميز مستخلص بذور العنب (GSE) الغني بالبروأنتوسيانيدينات، بخصائص مضادة للأكسدة قوية وقد يُقلل الضرر الناتج عن الكادميوم الذي يؤثر علي القدرة الإنجابية. تم تقسيم اربعون من ذكور الفئران البالغة إلى خمس مجموعات كل مجموعة تحتوى علي ٨ من الفئران : المجموعة الأولى (المجموعة الضابطة السالبة): اتبعت نظاماً غذائياً أساسياً فقط. المجموعة الثانية (مجموعة الضابطة الموجبة): اتبعت نظاماً غذائياً أساسياً وحُقت في التجويف الباطني بـ  $1 \text{ mg CdCl}_2 / \text{kg}$  من وزن الجسم لمدة ٨ أسابيع. المجموعات من ٣ إلى ٥: اتبعت نظاماً غذائياً أساسياً، وتناولت عن طريق الفم مستخلص بذور العنب GSE بجرعات ٢٠٠ أو ٤٠٠ أو ٨٠٠  $\text{mg/kg}$  من وزن الجسم، وحُقت بـ  $1 \text{ mg CdCl}_2 / \text{kg}$  من وزن الجسم. في نهاية فترة التجربة (شهرين)، تم تقدير عدة قياسات: هرمون التستوستيرون في السيرم، وكفاءة السائل المنوي. النتائج: أظهرت النتائج أن التعرض للكادميوم أدى الي إجهاد تأكسدي ملحوظ، وانخفاض ملحوظ في مستويات التستوستيرون، وتدهور ملحوظ في كفاءة الحيوانات المنوية، وقصراً في طول التيلوميرات. أظهرت النتائج الخاصة بمستخلص بذور العنب و المصابة بضعف في الخصية، وخاصةً بجرعتي ٤٠٠ و ٨٠٠ ملغم/كغم، تحسناً ملحوظاً في نشاط إنزيمات الكبد (ALT و AST)، ومستويات الدهون (TG) و TC و LDL-c و HDL-c)، ومستوى التستوستيرون الكلي في السيرم وعدد الحيوانات المنوية. كما أظهرت النتائج ان المجموعات التي تعاني من خلل وظيفي في الخصية وتغذت على نظام غذائي أساسي ومستخلص بذور العنب (٤٠٠، ٨٠٠ ملغم/كغم من وزن الجسم) زيادة كبيرة في جينات Casepase3 و STAR pcr و pcr علي تحليل شكل الحيوانات المنوية مقارنةً بالمجموعة الضابطة الموجبة(التي تعاني من خلل وظيفي في الخصية) وتغذت على نظام غذائي أساسي فقط. الخلاصة: مستخلص بذور العنب ربما يكون له تأثيرات وقائية ضد الخلل الوظيفي في الخصية الناجم عن الكادميوم، ويرجع ذلك إلى خصائصه المضادة للأكسدة وقدرته على تحسين نشاط التيلوميرات و جين اكسبريشين و الذي قد يكون مرتبطاً بالسموم البيئية.

الكلمات المفتاحية: التيلومير، المستخلص الإيثانولي لبذور العنب، الكادميوم، خلل وظيفي في الخصية، سمية الخصية و جين اكسبريشين..